

HIFEM FOR URINARY INCONTINENCE AND SEXUAL FUNCTION: 12-MONTH DATA

HIFEM PROCEDURE FOR IMPROVEMENT OF URINARY INCONTINENCE AND FEMALE SEXUAL FUNCTION: 12-MONTH MULTI-CENTER STUDY

Kimberly L. Evans¹; Joseph Berenholz²; Julene B. Samuels³
Andrea Pezzela⁴; Carolyn A. DeLucia⁵

1. Department of Cosmetic gynecology, Sugarland Medical Spa, Sugar Land, USA
2. Department of Obstetrics & Gynecology, Michigan Center for Women's Health, Farmington Hills, USA
3. Department of Plastic Surgery, Julene B. Samuels MD, FACS, Prospect, KY, USA
4. Department of Urogynecology, Southern Urogynecology, West Columbia, USA
5. Department of Obstetrics & Gynecology, Gynecological Solution, Hillsborough, NJ, USA


Published in Journal of Women's Health Care: February 2023, DOI: 10.35248/2167-0420.23.12.625

HIGHLIGHTS


- 31 females (30-62 years old) who showed a decreased interest in sexual activity accompanied by urinary incontinence, were enrolled in this study.

Results at 12 months:

- Urinary incontinence (ICIQ-UI) showed improvement by **68%**.
- Sexual dysfunction symptoms (FSFI) showed improvement by **51%**.
- Patients reported improved orgasmic response (**62%**).
- Significant increase was recorded in lubrication domain (**47%**).



ORGASM
IMPROVEMENT



LUBRICATION
INCREASE